

MIGRAINE & OTHER HEADACHES:

the role of the optometrist


Prof Bruce Evans

BSc (Hons) PhD FCOptom FAAO FEAOO FBCLA DipCLP DipOrth

Director of Research
Visiting Professor
Visiting Professor
Private practice

Institute of Optometry
City, University London
London South Bank University
Brentwood, Essex

© 1990-2019 Bruce Evans




Full handout from: www.bruce-evans.co.uk

for regular tweets on optometric research

2


DISCLOSURE

- Funding for lectures, KOL/product feedback, research:
 - Alcon, American Academy of Optometry (UK), Association of Optometrists, Birmingham Focus on Blindness, Black & Lizars, Butterworth-Heinemann (Elsevier), Central (L/OC) Fund, Cerium Visual Technologies, College of Optometrists, Coopervision, ESRC, General Optical Council, Hoya, International Institute for Colorimetry, Institute of Optometry, Iris Fund for Prevention of Blindness, Johnson & Johnson, Leightons, MRC, Norville, Optos, Paul Hamlyn Trust, Perceptive, Scrivens, Specsavers, Thomas Pocklington Trust.
 - Lecture content always my own
- Author of Pickwell's Binocular Vision Anomalies, editions 3-5
- i.O.O. Sales Ltd markets IFS Orthoptic Exercises, which the speaker designed, for which he receives a small royalty
- Community optometric practice in Brentwood, Essex



4

Role of the optometrist





Full handout from: www.bruce-evans.co.uk

for regular tweets on optometric research

6

Role of the optometrist




Full handout from: www.bruce-evans.co.uk

for regular tweets on optometric research

7

International Classification of Headache Disorders (ICHD-3, 2018)



8

4.	Other primary headache disorders	
4.1	Primary cough headache	A10.8.1 Headache attributed to travel in space
4.1.1	Probable primary cough headache	
4.2	Primary exercise headache	
4.2.1	Probable primary exercise headache	
4.3	Primary headache associated with sexual activity	
4.3.1	Probable primary headache associated with sexual activity	
4.4	Primary thunderclap headache	
4.5	Cold-stimulus headache	
4.5.1	Headache attributed to external application of a cold stimulus	
4.5.2	Headache attributed to ingestion or inhalation of a cold stimulus	
4.5.3	Probable cold-stimulus headache	
4.5.3.1	Headache probably attributed to external application of a cold stimulus	
4.5.3.2	Headache probably attributed to ingestion or inhalation of a cold stimulus	
4.6	External-pressure headache	
4.6.1	External-compression headache	
4.6.2	External-traction headache	
4.6.3	Probable external-pressure headache	
4.6.3.1	Probable external-compression headache	
4.6.3.2	Probable external-traction headache	
4.7	Primary stabbing headache	
4.7.1	Probable primary stabbing headache	...and many more

9



10

Prevalence in the real world

- Prevalence varies in different countries (Stovner et al., 2007)
- Marked difference in the two sexes (Lebedeva et al 2016)
- Marked effect of social groups & age (Lebedeva et al 2016)

Table 2. One-year crude prevalence of headache disorders according to ICHD-3 beta in blood donors, workers and students.

Diagnosis*	Male (n = 1775)			Female (n = 1349)			All (n = 3124)		
	Donors 33y (n = 523)	Workers 40y (n = 929)	Students 21y (n = 323)	Donors 33y (n = 484)	Workers 40y (n = 146)	Students 21y (n = 719)	Donors 33y (n = 1007)	Workers 40y (n = 1075)	Students 21y (n = 1042)
Migraine without aura	19 (3.6%)	41 (4.4%)	45 (13.9%)	74 (15.3%)	24 (16.4%)	202 (28.1%)	93 (9.2%)	65 (6.0%)	247 (23.7%)
Migraine with aura	4 (0.7%)	10 (1.1%)	2 (0.62%)	17 (3.5%)	3 (2%)	38 (5.3%)	21 (2.1%)	13 (1.2%)	40 (3.8%)
Chronic probable migraine	1 (0.2%)	0 (0%)	0 (0%)	3 (0.6%)	0 (0%)	1 (0.1%)	4 (0.4%)	0 (0%)	1 (0.1%)
Probable migraine	1 (0.2%)	1 (0.1%)	1 (0.3%)	0 (0%)	0 (0%)	9 (1.2%)	1 (0.1%)	1 (0.1%)	10 (0.9%)
Episodic TTH	307 (58.7%)	301 (32.4%)	250 (77.4%)	311 (64.3%)	94 (64.4%)	526 (73.2%)	618 (61.4%)	395 (36.7%)	776 (74.5%)
Chronic TTH	4 (0.8%)	1 (0.1%)	6 (1.9%)	7 (1.4%)	2 (1.4%)	26 (3.6%)	11 (1.1%)	3 (0.3%)	32 (3.1%)
Cluster headache	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	1 (0.1%)	0 (0%)	0 (0%)	1 (0.1%)
Medication-overuse headache (analgesics)	0 (0%)	4 (0.4%)	6 (1.9%)	8 (1.6%)	0 (0%)	26 (3.6%)	8 (0.8%)	4 (0.4%)	32 (3.1%)

*Diagnoses are not mutually exclusive.
TTH, tension-type headache; ICHD-3 beta, third edition beta of the International Classification of Headache Disorders.

11

Never ask a clinician about prevalence!

- In a headache clinic sample of 100, no secondary headaches and most were migraine Leone et al (1993)
- In an "emergency room", 26% were secondary headaches & the most common primary headache was migraine Munoz-Ceron et al. (2019)
- In both studies, no cases of ocular headache
- BUT, optometrists over-estimate prevalence of headaches from ocular origin

Headaches			13. Non-classifiable (8%)
Primary (91%)	Secondary (0%)	Secondary (cont.) (0%)	
1. Migraine (54%)	5. Head trauma	9. Non-cephalic infection	
2. Tension (30%)	6. Vascular	10. Metabolic disorder	
3. Cluster (7%)	7. Non-vascular intra-cranial disorder	11. Disorders of facial or cranial structures (inc. eye)	
4. Miscellaneous (0%)	8. Substances or their withdrawal	12. Neuralgias & deafferentation	

12

Role of the optometrist



Follow @BruceJWEvans

Full handout from: www.bruce-evans.co.uk for regular tweets on optometric research

13



14

ICHD-3 (cont.)

Headaches		
Primary	Secondary	Secondary (cont.)
1. Migraine	2. Headache due to trauma	1. Disorders of the cervical axis (e.g. occipital neuralgia)
2. Tension	3. Vascular	11. Disorders of facial or cranial structures (inc. eye)
3. Cluster	7. Non-vascular intra-cranial disorders	12. Neuralgias & deafferentation
4. Other primary headaches	8. Substances or their withdrawal	13. Other

Secondary

- 6. Vascular
 - 6.1 Ischaemic
 - 6.2 Haemorrhage (inc. sub-arachnoid)
 - 6.3 Vascular malformation (inc. aneurysm)
 - Arteritis
 - Hypertension
- 7. Non-vascular intra-cranial disorder
 - Intracranial hypertension
 - Intracranial hypotension
 - Intracranial infection
 - Intracranial inflamm. disease
 - Intrathecal injections
 - Intracranial neoplasm
 - Other intracranial disorder

- If sub-arachnoid haemorrhage:
 - Very severe (thunderclap), vomiting, photophobia
 - Bilateral

16


Differential diagnosis: the signs

1. Essentially, know the typical, refer the atypical
2. "first or worst" rule
3. SNOOP
 - a) **S**ystemic symptoms / signs / disease
 - b) **N**eurological Disease
 - c) **O**nset Sudden (thunderclap)
 - d) **O**nset after the age of forty
 - e) **P**attern change to the headaches



17

International Classification of Headache Disorders (ICHD-3, 2018)



21

Role of the optometrist



Follow @BruceJWEvans

Full handout from: www.bruce-evans.co.uk

for regular tweets on optometric research

22

Migraine: should I refer them all?

- * Migraine is third most prevalent disorder in the world (GBD2010)
- * Typical migraine does not need neuro-imaging (American Academy of Neurology, 1994; Detsky et al., 2006; Davies, 2018)
- * If diagnosis is obvious, no need for medical investigation
 - Letter of information to GP
- * If not very frequent, try analgesics first
 - See GP if not fully effective
- * Instruct to use diary
 - Identify any triggers → prevention



Full handout from: www.bruce-evans.co.uk

25

MIGRAINE WITHOUT AURA: at least 5 attacks which:

- last 4-72 hrs (untreated)
- have at least 2 of the following:
 - unilateral
 - pulsating
 - moderate/severe
 - aggravated by routine physical activity
- during HA at least 1 of the following
 - nausea and/or vomiting
 - photophobia and phonophobia

Is that it?


IHS (2018), ICHD-3

ICHD-3 code	Diagnosis
1.	Migraine
1.1	Migraine without aura
1.2	Migraine with aura
1.2.1	Migraine with typical aura
1.2.1.1	Typical aura with headache
1.2.1.2	Typical aura without headache
1.2.2	Migraine with brainstem aura
1.2.3	Hemiplegic migraine
1.2.3.1	Familial hemiplegic migraine (FHM)
1.2.3.1.1	Familial hemiplegic migraine type 1 (FHM1)
1.2.3.1.2	Familial hemiplegic migraine type 2 (FHM2)
1.2.3.1.3	Familial hemiplegic migraine type 3 (FHM3)
1.2.3.1.4	Familial hemiplegic migraine, other loci
1.2.3.2	Sporadic hemiplegic migraine (SHM)
1.2.4	Retinal migraine
1.3	Chronic migraine
1.4	Complications of migraine
1.4.1	Status migrainosus
1.4.2	Persistent aura without infarction
1.4.3	Migrainous infarction
1.4.4	Migraine aura-triggered seizure
1.5	Probable migraine
1.5.1	Probable migraine without aura
1.5.2	Probable migraine with aura
1.6	Episodic syndromes that may be associated with migraine
1.6.1	Recurrent gastrointestinal disturbance
1.6.1.1	Cyclical vomiting syndrome
1.6.1.2	Abdominal migraine
1.6.2	Benign paroxysmal vertigo
1.6.3	Benign paroxysmal torticollis

31

MIGRAINE WITH AURA: 2+ attacks with:


- fully reversible aura symptoms
 - e.g., vision, sensory, speech/language, motor, brainstem, retinal
- at least 3 of the following:
 - at least one aura symptom develops gradually over 5mins+ and/or
 - two or more symptoms occur in succession
 - each symptom lasts 5-60min
 - at least one aura symptom is unilateral
 - at least one aura symptom is positive
 - aura accompanied and/or followed with 60min by HA



IHS (2018), ICHD-3

32

Role of the optometrist



Full handout from: www.bruce-evans.co.uk

for regular tweets on optometric research

Follow @BruceJWEvans

34

Optometric correlates of migraine




- Subtle pupillary anomalies (Harle & Evans, 2004; Harle et al., 2005)
- Little evidence of visual field defects or increased risk of glaucoma (Harle & Evans, 2006a; Harle & Evans, 2005)
- Slightly higher prevalence of astigmatism and anisometropia (Harle & Evans, 2004; Harle & Evans, 2006c)
- Slightly higher prevalence of heterophoria and fixation disparity, but not usually a trigger (Harle & Evans, 2004; Harle & Evans, 2006b)
- The strongest visual correlate of migraine is pattern glare, which can be a migraine trigger (Harle & Evans, 2004; Harle et al., 2006)

39

Pattern glare

High prevalence in:

- Visual stress
- Migraine
- Photosensitive epilepsy
- Autism



DO NOT VIEW THIS IF YOU HAVE EPILEPSY OR MIGRAINE


46

Double-masked placebo-controlled trial of Precision Tints in people with migraine

INTRODUCTION: a small double-masked placebo-controlled trial of precision tinted lenses for migraine

RESULTS: Precision Tinted lenses reduce frequency of migraines for reasons that cannot be solely attributed to a placebo

STRENGTH: "suggestive"



Wilkins, Patel, Adjamian, Evans (2002)



Harle (2007): rare for migraineurs to need PTL

Speaker's opinion: evidence lacking, but probably extremely rare

49

PTL reduce cortical hyperactivation in migraine

- fMRI of 11 migraineurs & 11 non-headache controls
- Viewed visual stressful & non-stressful patterns through PTL (POT), control colour, grey
- Migraineurs & controls did not differ for non-stressful patterns
- Migraineurs had greater activation than controls with grey or control colours
 - Normalised with PTL



55

Do PTL alleviate visual triggers for migraine?




56

Why mention PTL?


Visual triggers for migraine are common

- Glare (62%)** (Debney, 1984)
 - Sun reflections (water, beach, snow, paper, chrome)
 - Bright lights
 - Windows [e.g., blinds]
- Flicker (53%)** (Debney, 1984)
 - Sun through trees/railings/etc
 - Stroboscopes (e.g., clubbing)
 - Faulty fluorescent lights [& non-faulty]
 - Television or cinema
 - Light in traffic tunnels
 - Flashlights or headlights
- Patterns** (Wilkins, 1995)
 - Carpets, escalators, shirts, text
- Rarely, neurologists refer particularly symptomatic pxs to optoms for colorimetry**



76

Role of the optometrist




Follow @BruceJWEvans

Full handout from: www.bruce-evans.co.uk

for regular tweets on optometric research

87

International Classification of Headache Disorders (ICHD-3, 2018)



88


Tension-type headache: >10 attacks which:

- last 30 mins - 7 days
- have at least 2 of the following:
 - bilateral
 - pressing/tightening (not pulsating)
 - mild/moderate (inhibits, not prohibits, daily activities)
 - not aggravated by routine physical activity
- both of the following
 - no nausea or vomiting
 - not both photophobia & phonophobia
- frequent, infrequent, and chronic varieties

IHS (2018), ICHD-3

91

International Classification of Headache Disorders (ICHD-3, 2018)



94

CLUSTER HA: 5+ attacks which:


- severe unilateral orbital, supraorbital and/or temporal pain lasting 15-180mins untreated
- HA associated with at least 1 of the following:
 - ipsilateral conjunctival injection and/or lacrimation
 - ipsilateral nasal congestion and/or running nose
 - ipsilateral eyelid oedema
 - ipsilateral forehead and facial sweating
 - ipsilateral miosis and/or ptosis
 - sense of restlessness and/or agitation
- frequency of attacks: 1 every other day to 8 per day

3.1	Cluster headache
3.1.1	Episodic cluster headache
3.1.2	Chronic cluster headache
3.2	Paroxysmal hemicrania
3.2.1	Episodic paroxysmal hemicrania
3.2.2	Chronic paroxysmal hemicrania
3.3	Short-lasting unilateral neuralgiform headache attacks
3.3.1	Short-lasting unilateral neuralgiform headache attacks with conjunctival injection and tearing (SUNCT)
3.3.1.1	Episodic SUNCT
3.3.1.2	Chronic SUNCT
3.3.2	Short-lasting unilateral neuralgiform headache attacks with cranial autonomic symptoms (SUNA)
3.3.2.1	Episodic SUNA
3.3.2.2	Chronic SUNA
3.4	Hemicrania continua
3.4.1	Hemicrania continua, remitting subtype
3.4.2	Hemicrania continua, unremitting subtype
3.5	Probable trigeminal autonomic cephalalgia
3.5.1	Probable cluster headache
3.5.2	Probable paroxysmal hemicrania
3.5.3	Probable short-lasting unilateral neuralgiform headache attacks
3.5.4	Probable hemicrania continua

IHS (2018), ICHD-3

96


International Classification of Headache Disorders (ICHD-3, 2018)



102

International Classification of Headache Disorders (ICHD-3)

- Compared with ICHD-2:
 - Relegated heterophoria to Appendix "because of insufficient evidence of their existence"



107

11.3 EYES: 11.3.2 Refractive errors

- un- or mis-corrected Rx
- absent on wakening, worse with prolonged relevant tasks
- evidence of causation
 - Temporal relation to Rx
 - Improves after Rx corrected
 - Aggravated by prolonged visual tasks
 - Improves when task discontinued

109

Asthenopia headache: received wisdom

- “Headache is the commonest symptom associated with eyestrain. This occurs in almost every possible variety”
- All “obscure headaches” should have eye exam before medical treatment. Rule out:
 - Environmental factors
 - Refractive error
 - Binocular vision anomaly
 - IOP, fields, discs, vessels (Duke-Elder, 1970)
- No form of headache is specific to eye-related disorders (Ball, 1982)

118

Does refractive error headache exist?

1a. Systematic review of homogenous RCTs
 1b. Individual RCT with good CI
 2a. Systematic review of homogenous cohort studies
 2b. Individual cohort study
 3a. Systematic review of case control studies
 3b. Individual case control study
 4. Case series
 5. Expert opinion

EBP is “the integration of best research evidence with clinical expertise and patient values.” (Sackett, 2000)

119

Role of the optometrist



Full handout from: www.bruce-evans.co.uk

for regular tweets on optometric research

Follow @BruceJWEvans

124

Headache diary

Enter every headache you experience in the table below. Impact ratings can be graded as: none, mild, severe, totally disabled
Intensity ratings can be graded as: mild, moderate, severe, very severe. Tick any of the "triggers" that describe your activities before the headache. "Hormonal" refers to certain times in women's monthly cycle; "flickering" refers to flickering fluorescent lights, discos, or lights flickering through trees; "patterns" refers to striped patterns. When you have completed the table, if any frequent triggers become apparent then try avoiding these. If visual stimuli (flickering, patterns, reading) are triggers then precision tinted lenses may help. At Cole Martin Trepaskis Optometrists we have a special instrument, the Intuitive Colorimeter, to test for these, please telephone for an appointment.

Date	Time	Impact	Intensity	Medicine Taken	Dosage	Time to Relief	Reading	Patterns	Flickering	Other triggers	Red wine	Other food	Others	Stress	Hormonal	Tiredness	Noise	Comments

Full handout from: www.bruce-evans.co.uk

127

Role of the optometrist

Full handout from: www.bruce-evans.co.uk


Follow @BruceJWEvans
for regular tweets on optometric research

137

Conclusions

- ☀️ Optometrists can diagnose commonplace headaches
 - 🔵 Know the normal: refer the abnormal asap
- ☀️ Refractive errors or binocular vision anomalies may be a trigger for headaches
- ☀️ Precision tinted lenses may help some people with headaches (migraine)
 - 🔵 These people usually know they have visual triggers

Full handout from: www.bruce-evans.co.uk



138

A personal perspective: Dr Optometry

- ☀️ In 2008 the Institute of Optometry launched a Doctor of Optometry degree in collaboration with London South Bank University
- ☀️ 5 year part time professional doctorate
 - 🔵 Year 1 has 13 taught days & 2 assignments
 - 🔵 Year 2 has 8 taught days & 2 assignments
 - 🔵 Years 3-5 are supervised doctoral research
 - 🔵 Research most likely to be clinical, in practice
- ☀️ "the ultimate qualification for UK optometrists"




141



Full handout from: www.bruce-evans.co.uk